

# Amine Boranes Dehydrogenation Mediated by an Unsymmetrical Iridium Pincer Hydride: (PCN) vs (PCP) Improved Catalytic Performance

Luconi L., Osipova E., Giambastiani G., Peruzzini M., Rossin A., Belkova N., Filippov O., Titova E., Pavlov A., Shubina E.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2018 American Chemical Society. The IrIII hydride (tBuPCN)IrHCl (1) containing the tridentate unsymmetrical pincer ligand tBuPCN- {tBuPCN(H) = 1-[3-[(di-*t*-rt-butylphosphino)methyl]phenyl]-1H-pyrazole} has been exploited as ammonia borane (NH<sub>3</sub>BH<sub>3</sub>, AB) and amine boranes dehydrogenation catalyst in THF solution at ambient temperature. 1 releases one H<sub>2</sub> equivalent per AB equivalent, with concomitant cyclic poly(aminoboranes) formation [B-(cyclotriborazanyl)-amine-borane (BCTB) and cyclotriborazane (CTB)] as the final "spent fuel". 1 has been found to have superior catalytic activity than its symmetrical analogue (tBuPCP)IrHCl, with recorded TOF values of 580 h<sup>-1</sup> (AB in THF) and 401 h<sup>-1</sup> (DMAB in toluene) at ambient temperature. The reaction has been analyzed experimentally through multinuclear [<sup>11</sup>B, <sup>31</sup>P{<sup>1</sup>H}, <sup>1</sup>H] NMR and IR spectroscopy, kinetic rate measurements, and kinetic isotope effect determination with deuterated AB isotopologues. The hydride/borohydride intermediate (tBuPCN)IrH(η<sup>2</sup>-BH<sub>4</sub>) (2) is the catalyst resting state formed during the dehydrogenation process; it is detected by a variable-temperature multinuclear NMR of the reaction course (in the 190–323 K range). A DFT modeling of the reaction mechanism using DMAB as substrate has been performed with the geometry optimization in toluene at the M06 level of theory. The combination of the kinetic and computational data reveals that a simultaneous B-H/N-H activation occurs in the presence of 1, after the preliminary amine borane coordination to the metal center.

<http://dx.doi.org/10.1021/acs.organomet.8b00488>

---

## References

- [1] Rossin, A.; Peruzzini, M. Ammonia-Borane and Amine-Borane Dehydrogenation Mediated by Complex Metal Hydrides. *Chem. Rev.* 2016, 116, 8848–8872, 10.1021/acs.chemrev.6b00043
- [2] Yadav, M.; Xu, Q. Liquid-Phase Chemical Hydrogen Storage Materials. *Energy Environ. Sci.* 2012, 5, 9698–9725, 10.1039/c2ee22937d
- [3] Demirci, U. B.; Miele, P. Chemical Hydrogen Storage: 'Material' Gravimetric Capacity Versus 'System' Gravimetric Capacity. *Energy Environ. Sci.* 2011, 4, 3334–3341, 10.1039/c1ee01612a
- [4] Jiang, H.-L.; Xu, Q. Catalytic Hydrolysis of Ammonia Borane for Chemical Hydrogen Storage. *Catal. Today* 2011, 170, 56–63, 10.1016/j.cattod.2010.09.019
- [5] Sanyal, U.; Demirci, U. B.; Jagirdar, B. R.; Miele, P. Hydrolysis of Ammonia Borane as a Hydrogen Source: Fundamental Issues and Potential Solutions Towards Implementation. *ChemSusChem* 2011, 4, 1731–1739, 10.1002/cssc.201100318

- [6] Hamilton, C. W.; Baker, R. T.; Staubitz, A.; Manners, I. B-N Compounds for Chemical Hydrogen Storage. *Chem. Soc. Rev.* 2009, 38, 279-293, 10.1039/B800312M
- [7] Hügler, T.; Hartl, M.; Lentz, D. The Route to a Feasible Hydrogen-Storage Material: MOFs Versus Ammonia Borane. *Chem.-Eur. J.* 2011, 17, 10184-10207, 10.1002/chem.201003364
- [8] Staubitz, A.; Robertson, A. P. M.; Manners, I. Ammonia-Borane and Related Compounds as Dihydrogen Sources. *Chem. Rev.* 2010, 110, 4079-4124, 10.1021/cr100088b
- [9] Smythe, N. C.; Gordon, J. C. Ammonia Borane as a Hydrogen Carrier: Dehydrogenation and Regeneration. *Eur. J. Inorg. Chem.* 2010, 2010, 509-521, 10.1002/ejic.200900932
- [10] Stephens, F. H.; Pons, V.; Baker, R. T. Ammonia-Borane: the Hydrogen Source par Excellence?. *Dalton Trans.* 2007, 2613-2626, 10.1039/B703053C
- [11] Marder, T. B. Will We Soon Be Fueling our Automobiles with Ammonia-Borane?. *Angew. Chem., Int. Ed.* 2007, 46, 8116-8118, 10.1002/anie.200703150
- [12] Luconi, L.; Lyubov, D. M.; Rossin, A.; Glukhova, T. A.; Cherkasov, A. V.; Tuci, G.; Fukin, G. K.; Trifonov, A. A.; Giambastiani, G. Organolanthanide Complexes Supported by Thiazole-Containing Amidopyridinate Ligands: Synthesis, Characterization, and Catalytic Activity in Isoprene Polymerization. *Organometallics* 2014, 33, 7125-7134, 10.1021/om500918y
- [13] Wallis, C. J.; Dyer, H.; Vendier, L.; Alcaraz, G.; Sabo-Etienne, S. Dehydrogenation of Diamine-Monoboranes to Cyclic Diaminoboranes: Efficient Ruthenium-Catalyzed Dehydrogenative Cyclization. *Angew. Chem., Int. Ed.* 2012, 51, 3646-3648, 10.1002/anie.201108874
- [14] Conley, B. L.; Guess, D.; Williams, T. J. A Robust, Air-Stable, Reusable Ruthenium Catalyst for Dehydrogenation of Ammonia Borane. *J. Am. Chem. Soc.* 2011, 133, 14212-14215, 10.1021/ja2058154
- [15] Alcaraz, G.; Chaplin, A. B.; Stevens, C. J.; Clot, E.; Vendier, L.; Weller, A. S.; Sabo-Etienne, S. Ruthenium, Rhodium, and Iridium bis( $\sigma$ -B-H) Diisopropylaminoborane Complexes. *Organometallics* 2010, 29, 5591-5595, 10.1021/om1004995
- [16] Conley, B. L.; Williams, T. J. Dehydrogenation of Ammonia-Borane by Shvo's Catalyst. *Chem. Commun.* 2010, 46, 4815-4817, 10.1039/c003157g
- [17] Käß, M.; Friedrich, A.; Drees, M.; Schneider, S. Ruthenium Complexes with Cooperative PNP Ligands: Bifunctional Catalysts for the Dehydrogenation of Ammonia-Borane. *Angew. Chem., Int. Ed.* 2009, 48, 905-907, 10.1002/anie.200805108
- [18] Blaquiere, N.; Diallo-Garcia, S.; Gorelsky, S. I.; Black, D. A.; Fagnou, K. Ruthenium-Catalyzed Dehydrogenation of Ammonia Boranes. *J. Am. Chem. Soc.* 2008, 130, 14034-14035, 10.1021/ja804235t
- [19] Sewell, L. J.; Lloyd-Jones, G. C.; Weller, A. S. Development of a Generic Mechanism for the Dehydrocoupling of Amine-Boranes: A Stoichiometric, Catalytic, and Kinetic Study of HB-NMeH Using the [Rh(PCy)] Fragment. *J. Am. Chem. Soc.* 2012, 134, 3598-3610, 10.1021/ja2112965
- [20] Tang, C. Y.; Thompson, A. L.; Aldridge, S. Rhodium and Iridium Aminoborane Complexes: Coordination Chemistry of BN Alkene Analogues. *Angew. Chem., Int. Ed.* 2010, 49, 921-925, 10.1002/anie.200906171
- [21] Dallanegra, R.; Chaplin, A. B.; Weller, A. S. Bis( $\sigma$ -Amine-Borane) Complexes: An Unusual Binding Mode at a Transition-Metal Center. *Angew. Chem., Int. Ed.* 2009, 48, 6875-6878, 10.1002/anie.200903121
- [22] Douglas, T. M.; Chaplin, A. B.; Weller, A. S.; Yang, X.; Hall, M. B. Monomeric and Oligomeric Amine-Borane  $\sigma$ -Complexes of Rhodium. Intermediates in the Catalytic Dehydrogenation of Amine-Boranes. *J. Am. Chem. Soc.* 2009, 131, 15440-15456, 10.1021/ja906070r
- [23] Douglas, T. M.; Chaplin, A. B.; Weller, A. S. Amine-Borane  $\sigma$ -Complexes of Rhodium. Relevance to the Catalytic Dehydrogenation of Amine-Boranes. *J. Am. Chem. Soc.* 2008, 130, 14432-14433, 10.1021/ja806582n
- [24] Chen, Y.; Fulton, J. L.; Linehan, J. C.; Autrey, T. In Situ XAFS and NMR Study of Rhodium-Catalyzed Dehydrogenation of Dimethylamine Borane. *J. Am. Chem. Soc.* 2005, 127, 3254-3255, 10.1021/ja0437050
- [25] Johnson, H. C.; Robertson, A. P. M.; Chaplin, A. B.; Sewell, L. J.; Thompson, A. L.; Haddow, M. F.; Manners, I. A.; Weller, A. S. Catching the First Oligomerization Event in the Catalytic Formation of Polyaminoboranes: HB-NMeHBH-NMeH Bound to Iridium. *J. Am. Chem. Soc.* 2011, 133, 11076-11079, 10.1021/ja2040738
- [26] Rossin, A.; Caporali, M.; Gonsalvi, L.; Guerri, A.; Lledós, A.; Peruzzini, M.; Zanobini, F. Selective B-H versus N-H Bond Activation in Ammonia Borane by [Ir(dppm)]OTf. *Eur. J. Inorg. Chem.* 2009, 2009, 3055-3059, 10.1002/ejic.200900465
- [27] Paul, A.; Musgrave, C. B. Catalyzed Dehydrogenation of Ammonia-Borane by Iridium Dihydrogen Pincer Complex Differs from Ethane Dehydrogenation. *Angew. Chem., Int. Ed.* 2007, 46, 8153-8156, 10.1002/anie.200702886
- [28] Denney, M. C.; Pons, V.; Hebden, T. J.; Heinekey, M.; Goldberg, K. I. Efficient Catalysis of Ammonia Borane Dehydrogenation. *J. Am. Chem. Soc.* 2006, 128, 12048-12049, 10.1021/ja062419g
- [29] Szabo, K. J.; Wendt, O. F. Pincer and Pincer-Type Complexes: Application in Organic Synthesis and Catalysis; Wiley-VCH: Weinheim, Germany, 2014.
- [30] Van Koten, G.; Milstein, D. Organometallic Pincer Chemistry; Springer: Heidelberg, 2013; Vol. 40.

- [31] Benito-Garagorri, D.; Kirchner, K. Modularly Designed Transition Metal PNP and PCP Pincer Complexes Based on Aminophosphines: Synthesis and Catalytic Applications. *Acc. Chem. Res.* 2008, 41, 201-213, 10.1021/ar700129q
- [32] van der Boom, M. E.; Milstein, D. Cyclometalated Phosphine-Based Pincer Complexes: Mechanistic Insight in Catalysis, Coordination, and Bond Activation. *Chem. Rev.* 2003, 103, 1759-1792, 10.1021/cr960118r
- [33] Singleton, J. T. The Uses of Pincer Complexes in Organic Synthesis. *Tetrahedron* 2003, 59, 1837-1857, 10.1016/S0040-4020(02)01511-9
- [34] Moulton, C. J.; Shaw, B. L. Transition Metal-Carbon Bonds. Part XIII. Complexes of Nickel, Palladium, Platinum, Rhodium and Iridium with the Tridentate Ligand 2,6-Bis[(di- *t*-Butylphosphino)Methyl]Phenyl. *J. Chem. Soc., Dalton Trans.* 1976, 1020-1024, 10.1039/DT9760001020
- [35] Roddick, D. M. Tuning of PCP Pincer Ligand Electronic and Steric Properties. *Top. Organomet. Chem.* 2013, 40, 49-88, 10.1007/978-3-642-31081-2-3
- [36] Kundu, S.; Choliy, Y.; Zhuo, G.; Ahuja, R.; Emge, T. J.; Warmuth, R.; Brookhart, M.; Krogh-Jespersen, K.; Goldman, A. S. Rational Design and Synthesis of Highly Active Pincer-Iridium Catalysts for Alkane Dehydrogenation. *Organometallics* 2009, 28, 5432-5444, 10.1021/om900568f
- [37] Leis, W.; Mayer, H. A.; Kaska, W. C. Cycloheptatrienyl, Alkyl and Aryl PCP-Pincer Complexes: Ligand Backbone Effects and Metal Reactivity. *Coord. Chem. Rev.* 2008, 252, 1787-1797, 10.1016/j.ccr.2008.02.002
- [38] Jensen, C. M. Iridium PCP Pincer Complexes: Highly Active and Robust Catalysts for Novel Homogeneous Aliphatic Dehydrogenations. *Chem. Commun.* 1999, 2443-2449, 10.1039/a903573g
- [39] Schneider, S.; Meiners, J.; Askevold, B. Cooperative Aliphatic PNP Amido Pincer Ligands-Versatile Building Blocks for Coordination Chemistry and Catalysis. *Eur. J. Inorg. Chem.* 2012, 2012, 412-429, 10.1002/ejic.201100880
- [40] Milstein, D. Discovery of Environmentally Benign Catalytic Reactions of Alcohols Catalyzed by Pyridine-Based Pincer Ru Complexes, Based on Metal-Ligand Cooperation. *Top. Catal.* 2010, 53, 915-923, 10.1007/s11244-01-9523-7
- [41] Timpa, S. D.; Pell, C.; Zhou, J.; Ozerov, O. V. Carbon-Halide Oxidative Addition and Carbon-Carbon Reductive Elimination at a (PNP)Rh Center. *Organometallics* 2014, 33, 5254-5262, 10.1021/om500577w
- [42] Jonasson, K. J.; Wendt, O. F. Synthesis and Characterization of a Family of POCOP Pincer Complexes with Nickel: Reactivity Towards CO and Phenylacetylene. *Chem.-Eur. J.* 2014, 20, 11894-11902, 10.1002/chem.201403246
- [43] Lao, D. B.; Owens, A. C. E.; Heinekey, D. M.; Goldberg, K. I. Partial Deoxygenation of Glycerol Catalyzed by Iridium Pincer Complexes. *ACS Catal.* 2013, 3, 2391-2396, 10.1021/cs400551g
- [44] Pandarus, V.; Zargarian, D. New Pincer-Type Diphosphinito (POCOP) Complexes of Ni And Ni. *Chem. Commun.* 2007, 978-980, 10.1039/B613812H
- [45] Hebden, T. J.; St. John, A. J.; Gusev, D. G.; Kaminsky, W.; Goldberg, K. I.; Heinekey, D. M. Preparation of a Dihydrogen Complex of Cobalt. *Angew. Chem., Int. Ed.* 2011, 50, 1873-1876, 10.1002/anie.201005281
- [46] Hao, X.-Q.; Huang, J.-J.; Wang, T.; Lv, J.; Gong, J.-F.; Song, M.-P. PCN Pincer Palladium(II) Complex Catalyzed Enantioselective Hydrophosphination of Enones: Synthesis of Pyridine-Functionalized Chiral Phosphine Oxides as NCO Pincer Preligands. *J. Org. Chem.* 2014, 79, 9512-9530, 10.1021/jo5015307
- [47] Khake, S. M.; Soni, V.; Gonnade, R. G.; Punji, B. Design and Development of POCN-Pincer Palladium Catalysts for C-H Bond Arylation of Azoles with Aryl Iodides. *Dalton Trans.* 2014, 43, 16084-16096, 10.1039/C4DT01547A
- [48] Herbert, D. E.; Ozerov, O. V. Binuclear Palladium Complexes Supported by Bridged Pincer Ligands. *Organometallics* 2011, 30, 6641-6654, 10.1021/om2008056
- [49] Moreno, I.; SanMartin, R.; Inés, B.; Herrero, M. T.; Domínguez, E. Recent Advances in the Use of Unsymmetrical Palladium Pincer Complexes. *Curr. Org. Chem.* 2009, 13, 878-895, 10.2174/138527209788452144
- [50] Poverenov, E.; Leitus, G.; Shimon, L. J. W.; Milstein, D. C-Metalated Diazoalkane Complexes of Platinum Based on PCP- and PCN-Type Ligands. *Organometallics* 2005, 24, 5937-5944, 10.1021/om050637x
- [51] Bailey, W.; Luconi, L.; Rossin, A.; Yakhvarov, D.; Flowers, S. E.; Kaminsky, W.; Kemp, R. A.; Giambastiani, G.; Goldberg, K. I. Pyrazole-Based PCN Pincer Complexes Of Palladium(II): Mono- and Dinuclear Hydroxide Complexes and Ligand Rollover C-H Activation. *Organometallics* 2015, 34, 3998-4010, 10.1021/acs.organomet.5b00355
- [52] Zhang, X.; Suzuki, S.; Kozaki, M.; Okada, K. NCN Pincer-Pt Complexes Coordinated by (Nitronyl Nitroxide)-2-ide Radical Anion. *J. Am. Chem. Soc.* 2012, 134, 17866-17868, 10.1021/ja308103g
- [53] Gong, J.-F.; Zhang, Y.-H.; Song, M.-P.; Xu, C. New PCN and PCP Pincer Palladium(II) Complexes: Convenient Synthesis via Facile One-Pot Phosphorylation/Palladation Reaction and Structural Characterization. *Organometallics* 2007, 26, 6487-6492, 10.1021/om7008364
- [54] Titova, E. M.; Osipova, E. S.; Pavlov, A. A.; Filippov, O. A.; Safronov, S. V.; Shubina, E. S.; Belkova, N. V. Mechanism of Dimethylamine-Borane Dehydrogenation Catalyzed by an Iridium(III) PCP-Pincer Complex. *ACS Catal.* 2017, 7, 2325-2333, 10.1021/acscatal.6b03207

- [55] Todisco, S.; Luconi, L.; Giambastiani, G.; Rossin, A.; Peruzzini, M.; Golub, I. E.; Filippov, O. A.; Belkova, N. V.; Shubina, E. S. Ammonia Borane Dehydrogenation Catalyzed by ( $\kappa$ -EP)Co(H) [EP = E(CHCHPPh); E = N, P] and H Evolution from Their Interaction with NH Acids. *Inorg. Chem.* 2017, 56, 4296-4307, 10.1021/acs.inorgchem.6b02673
- [56] Luconi, L.; Rossin, A.; Motta, A.; Tuci, G.; Giambastiani, G. Group IV Organometallic Compounds Based on Dianionic "Pincer" Ligands: Synthesis, Characterization, and Catalytic Activity in Intramolecular Hydroamination Reactions. *Chem.-Eur. J.* 2013, 19, 4906-4921, 10.1002/chem.201203644
- [57] Rossin, A.; Peruzzini, M.; Zanobini, F. Nickel(II) Hydride and Fluoride Pincer Complexes and their Reactivity with Lewis Acids BX<sub>3</sub>L (X = H, L = thf; X = F, L = EtO). *Dalton Trans.* 2011, 40, 4447-4452, 10.1039/c0dt01836h
- [58] Luconi, L.; Gafurov, Z.; Rossin, A.; Tuci, G.; Sinyashin, O.; Yakhvarov, D.; Giambastiani, G. Palladium(II) Pyrazolyl-Pyridyl Complexes Containing a Sterically Hindered N-Heterocyclic Carbene Moiety for the Suzuki-Miyaura Cross-Coupling Reaction. *Inorg. Chim. Acta* 2018, 470, 100-105, 10.1016/j.ica.2017.03.026
- [59] Herde, J. L.; Lambert, J. C.; Senoff, C. V.; Cushing, M. A. Cyclooctene and 1,5-Cyclooctadiene Complexes of Iridium(I). *Inorg. Synth.* 1982, 15, 19-20, 10.1002/9780470132463.ch5
- [60] Hu, M. G.; Van Paasschen, J. M.; Geanangel, R. A. New Synthetic Approaches to Ammonia-Borane and its Deuterated Derivatives. *J. Inorg. Nucl. Chem.* 1977, 39, 2147-2150, 10.1016/0022-1902(77)80383-7
- [61] CrysAlis CCD 1.171.31.2 (release 07-07-2006, CrysAlis171.NET); Rigaku Oxford Diffraction Ltd.: The Woodlands, TX, 2006.
- [62] CrysAlis RED 1.171.31.2 (release 07-07-2006, CrysAlis171.NET); Rigaku Oxford Diffraction Ltd.: The Woodlands, TX, 2006.
- [63] Altomare, A.; Burla, M. C.; Camalli, M.; Cascarano, G. L.; Giacovazzo, C.; Guagliardi, A.; Moliterni, A. G. G.; Polidori, G.; Spagna, R. SIR97: a New Tool for Crystal Structure Determination and Refinement. *J. Appl. Crystallogr.* 1999, 32, 115-119, 10.1107/S0021889898007717
- [64] Sheldrick, G. M. SHELXTL Version 2014/7; University of Göttingen: Göttingen, Germany, 2014.
- [65] Spek, A. L. Structure Validation in Chemical Crystallography. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* 2009, 65, 148-155, 10.1107/S090744490804362X
- [66] Nardelli, M. Parst: A System of Fortran Routines for Calculating Molecular Structure Parameters from Results of Crystal Structure Analyses. *Comput. Chem.* 1983, 7, 95-98, 10.1016/0097-8485(83)85001-3
- [67] Farrugia, L. J. ORTEP-3 for Windows-a Version of ORTEP-III with a Graphical User Interface (GUI). *J. Appl. Crystallogr.* 1997, 30, 565, 10.1107/S0021889897003117
- [68] Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, J. A., Jr.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; Pople, J. A. Gaussian 09; Gaussian, Inc.: Wallingford CT, 2009.
- [69] Zhao, Y.; Truhlar, D. G. The M06 Suite of Density Functionals for Main Group Thermochemistry, Thermochemical Kinetics, Noncovalent Interactions, Excited States, and Transition Elements: Two New Functionals and Systematic Testing of Four M06-Class Functionals and 12 Other Functionals. *Theor. Chem. Acc.* 2008, 120, 215-241, 10.1007/s00214-007-0310-x
- [70] Andrae, D.; Häußermann, U.; Dolg, M.; Stoll, H.; Preuß, H. Energy-Adjusted Ab initio Pseudopotentials for the Second and Third Row Transition Elements. *Theoret. Chim. Acta* 1990, 77, 123-141, 10.1007/BF01114537
- [71] Hausermann, U.; Dolg, M.; Stoll, H.; Preuss, H.; Schwerdtfeger, P.; Pitzer, R. M. Accuracy of Energy-Adjusted Quasirelativistic Ab Initio Pseudopotentials. *Mol. Phys.* 1993, 78, 1211-1224, 10.1080/00268979300100801
- [72] Kuchle, W.; Dolg, M.; Stoll, H.; Preuss, H. Energy-Adjusted Pseudopotentials for the Actinides. Parameter Sets and Test Calculations for Thorium and Thorium Monoxide. *J. Chem. Phys.* 1994, 100, 7535-7542, 10.1063/1.466847
- [73] Leininger, T.; Nicklass, A.; Stoll, H.; Dolg, M.; Schwerdtfeger, P. The Accuracy of the Pseudopotential Approximation. II. A Comparison of Various Core Sizes for Indium Pseudopotentials in Calculations for Spectroscopic Constants of InH, InF, and InCl. *J. Chem. Phys.* 1996, 105, 1052-1059, 10.1063/1.471950
- [74] Ehlers, A. W.; Dapprich, S.; Gobbi, A.; Höllwarth, A.; Jonas, V.; Köhler, K. F.; Stegmann, R.; Veldkamp, A.; Frenking, G.; Bohme, M. A Set of f-Polarization Functions for Pseudo-Potential Basis Sets of the Transition Metals Sc-Cu, Y-Ag and La-Au. *Chem. Phys. Lett.* 1993, 208, 111-114, 10.1016/0009-2614(93)80086-5

- [75] Clark, T. C. J.; Spitznagel, G. W.; Schleyer, P. V. R.; Chandrasekhar, J. Efficient Diffuse Function-Augmented Basis Sets for Anion Calculations. III. The 3-21+G Basis Set for First-Row Elements, Li-F. *J. Comput. Chem.* 1983, 4, 294-301, 10.1002/jcc.540040303
- [76] Francl, M. M. P. W. J.; Hehre, W. J.; Binkley, J. S.; DeFrees, D. J.; Pople, J. A.; Gordon, M. S.; Pietro, W. J. Self-Consistent Molecular Orbital Methods. XXIII. A Polarization-Type Basis Set for Second-Row Elements. *J. Chem. Phys.* 1982, 77, 3654-3665, 10.1063/1.444267
- [77] Hariharan, P. C. P. J. A.; Pople, J. A. The Influence of Polarization Functions on Molecular Orbital Hydrogenation Energies. *Theoret. Chim. Acta* 1973, 28, 213-222, 10.1007/BF00533485
- [78] Hehre, W. J. D. R.; Pople, J. A.; Ditchfield, R. Self-Consistent Molecular Orbital Methods. XII. Further Extensions of Gaussian-Type Basis Sets for Use in Molecular Orbital Studies of Organic Molecules. *J. Chem. Phys.* 1972, 56, 2257-2261, 10.1063/1.1677527
- [79] Marenich, A. V. C. C. J.; Truhlar, D. G.; Cramer, C. J. Universal Solvation Model Based on Solute Electron Density and on a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions. *J. Phys. Chem. B* 2009, 113, 6378-6396, 10.1021/jp810292n
- [80] Fritsch, J. Z. G.; Zundel, G. Influence of the Polarity of the Environment on Easily Polarizable OH - -N = O - -HN Hydrogen Bonds. *J. Phys. Chem.* 1981, 85, 556-561, 10.1021/j150605a020
- [81] Fukui, K. The Path of Chemical Reactions-the IRC Approach. *Acc. Chem. Res.* 1981, 14, 363-406, 10.1021/ar00072a001
- [82] Hratchian, H. P.; Schlegel, H. B. In *Theory and Applications of Computational Chemistry*; Frenking, G., Kim, K. S., Scuseria, G. E., Eds.; Elsevier: Amsterdam, 2005.
- [83] Titova, E. M.; Silantyev, G. A.; Filippov, O. A.; Gulyaeva, E. S.; Gutsul, E. I.; Dolgushin, F. M.; Belkova, N. V. PCP Pincer Iridium Chemistry-Coordination of Pyridines to [(PCP)IrH(Cl)]. *Eur. J. Inorg. Chem.* 2016, 2016, 56-63, 10.1002/ejic.201501083
- [84] Punji, B.; Emge, T. J.; Goldman, A. S. A Highly Stable Adamantyl-Substituted Pincer-Ligated Iridium Catalyst for Alkane Dehydrogenation. *Organometallics* 2010, 29, 2702-2709, 10.1021/om100145q
- [85] Mucha, N. T.; Waterman, R. Iridium Pincer Catalysts for Silane Dehydrocoupling: Ligand Effects on Selectivity and Activity. *Organometallics* 2015, 34, 3865-3872, 10.1021/acs.organomet.5b00486
- [86] Wang, J. S.; Geanangel, R. A. B NMR Studies of the Thermal Decomposition of Ammonia-Borane in Solution. *Inorg. Chim. Acta* 1988, 148, 185-190, 10.1016/S0020-1693(00)87499-X
- [87] Shaw, W. J.; Linehan, J. C.; Szymczak, N. K.; Heldebrant, D. J.; Yonker, C.; Camaioni, D. M.; Baker, R. T.; Autrey, T. In Situ Multinuclear NMR Spectroscopic Studies of the Thermal Decomposition of Ammonia Borane in Solution. *Angew. Chem., Int. Ed.* 2008, 47, 7493-7496, 10.1002/anie.200802100
- [88] Lingam, H. K.; Wang, C.; Gallucci, J. C.; Chen, X.; Shore, S. G. New Syntheses and Structural Characterization of NHBHCl and (BHNH) and Thermal Decomposition Behavior of NHBHCl. *Inorg. Chem.* 2012, 51, 13430-13436, 10.1021/ic302308v
- [89] Socrates, G. *Infrared and Raman Characteristic Group Frequencies: Tables and Charts*; John Wiley & Sons: New York, 2013.
- [90] Kumar, A.; Johnson, H. C.; Hooper, T. N.; Weller, A. S.; Algarra, A. G.; MacGregor, S. A. Multiple Metal-Bound Oligomers from Ir-Catalysed Dehydropolymerisation of HB·NH as Probed by Experiment and Computation. *Chem. Sci.* 2014, 5, 2546-2533, 10.1039/C4SC00735B
- [91] Gilbert, T. M. Tests of the MP2Model and Various DFT Models In Predicting the Structures and B-N Bond Dissociation Energies of Amine-Boranes (XC)HB-N(CH)H (X = H, F; m = 0-3; n = 0-3): Poor Performance of the B3LYP Approach for Dative B-N Bonds. *J. Phys. Chem. A* 2004, 108, 2550-2554, 10.1021/jp031242z